

(AS online)

***** Welcome to STN International *****

NEWS 1 Feb 2 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Dec 17 Expanded CAplus Coverage of US, Japanese, WIPO,
EPO, and German patents
NEWS 3 Feb 1 Addition of Machine-Translated Abstracts to CAplus
NEWS 4 Feb 2 STEPEC BOND SEARCH PROBLEM FIXED WITH STN EXPRESS 5.0C
NEWS 5 Feb 14 Homology Searching for Nucleotide Sequences in DGENE
now available!
NEWS 6 Feb 16 BIOTECHNOBASE NOW ON STN
NEWS 7 Feb 23 New Database Producer Clusters Now Available on STN
NEWS 8 Feb 23 Structure Search Limits Increased in REGISTRY,
ZEPIDESTRY, and CASREACT
NEWS 9 Feb 23 Patent Information Now Searchable in CAOLD
NEWS 10 Mar 1 New IMSDIRECTORY Provides Pharma Company Details
NEWS 11 Mar 20 INPADOC: PRODUCER WARNING ABOUT DATA DELAYS
NEWS 12 Mar 22 NEW FEATURES IN INPADOC - RANGE SEARCHING AND NEW
SLI/UPDATE SEARCH FIELD

NEWS EXPRESS FREE UPGRADE 5.0C NOW AVAILABLE
NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

***** STN Columbus *****

FILE 'HOME' ENTERED AT 15:11:03 ON 23 MAR 2000

=> file chemistry

FILE 'APILIT2' ACCESS NOT AUTHORIZED

FILE 'PAPERCHEM' ACCESS NOT AUTHORIZED

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'AGRICOLA' ENTERED AT 15:11:42 ON 23 MAR 2000

FILE 'ALUMINIUM' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Cambridge Scientific Abstracts (CSA)

FILE 'ANABSTR' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 THE ROYAL SOCIETY OF CHEMISTRY (RSC)

FILE 'APILIT' ENTERED AT 15:11:42 ON 23 MAR 2000

COPYRIGHT (C) 2000 Elsevier Engineering Information, Inc. (Elsevier)

FILE 'BIOCOMMERCE' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 BioCommerce Data Ltd. Richmond Surrey, United Kingdom.
All rights reserved

FILE 'BIOTECHNO' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Elsevier Science B.V., Amsterdam. All rights reserved.

FILE 'CABA' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 CAB INTERNATIONAL (CABI)

FILE 'CAOLD' ENTERED AT 15:11:42 ON 23 MAR 2000
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2000 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'CAPLUS' ENTERED AT 15:11:42 ON 23 MAR 2000
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2000 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'CENB' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 THE ROYAL SOCIETY OF CHEMISTRY (RSC)

FILE 'CEABA' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 DECHEMA eV

FILE 'CEN' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 American Chemical Society (ACS)

FILE 'CERAB' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Cambridge Scientific Abstracts (CSA)

FILE 'CIN' ENTERED AT 15:11:42 ON 23 MAR 2000
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2000 American Chemical Society (ACS)

FILE 'COMPENDEX' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 ENGINEERING INFORMATION, INC. (EI)

FILE 'CONFSCI' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Cambridge Scientific Abstracts (CSA)

FILE 'GENBANK' ENTERED AT 15:11:42 ON 23 MAR 2000

FILE 'INSPEC' ENTERED AT 15:11:42 ON 23 MAR 2000
Compiled and produced by the IEE in association with FIZ KARLSRUHE
COPYRIGHT 2000 (C) INSTITUTION OF ELECTRICAL ENGINEERS (IEE)

FILE 'INSPHYS' ENTERED AT 15:11:42 ON 23 MAR 2000
Compiled and produced by the IEE in association with FIZ KARLSRUHE
COPYRIGHT 2000 (C) INSTITUTION OF ELECTRICAL ENGINEERS (IEE)

FILE 'INVESTEXT' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Thomson Financial Services, Inc. (TFS)

FILE 'IPA' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 American Society of Hospital Pharmacists (ASHP)

FILE 'JICST-EPLUS' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Japan Science and Technology Corporation (JST)

FILE 'KKF' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (c) 2000 Deutsches Kunststoff-Institut (DKI)

FILE 'KOSMET' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 International Federation of the Societies of Cosmetics Chemists

FILE 'METADEX' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (c) 2000 Cambridge Scientific Abstracts (CSA)

FILE 'NAPRALERT' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Board of Trustees of the University of Illinois,
University of Illinois at Chicago.

FILE 'NIOSHTIC' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 U.S. Secretary of Commerce on Behalf of the U.S.
Government

FILE 'NTIS' ENTERED AT 15:11:42 ON 23 MAR 2000
Compiled and distributed by the NTIS, U.S. Department of Commerce.
It contains copyrighted material.
All rights reserved. (2000)

FILE 'PAPERCHEM2' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Institute of Paper Science and Technology (IPST)

FILE 'PROMT' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Gale Group. All rights reserved.

FILE 'RAPRA' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 RAPRA Technology Ltd.

FILE 'RUSSCI' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Andrigal Ltd.

FILE 'SCISEARCH' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 Institute for Scientific Information (ISI) (R)

FILE 'TULSA' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 The University of Tulsa (UTULSA)

FILE 'TULSA2' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 The University of Tulsa (UTULSA)

FILE 'USAN' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 U.S. Pharmaceutical Convention, Inc. (USPC)

FILE 'VTB' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (c) 2000 BAYER AG/BASF Aktiengesellschaft/DECHEMA eV

FILE 'WSCA' ENTERED AT 15:11:42 ON 23 MAR 2000
COPYRIGHT (C) 2000 PAINT RESEARCH

=> s tocopherol

21 FILES SEARCHED...

28 FILES SEARCHED...

L1 43093 TOCOPHEROL

=> s l1 and separat?

18 FILES SEARCHED...

L2 2438 L1 AND SEPARAT?

=> s l2 and esterify?

37 FILES SEARCHED...

L: 6 L2 AND ESTERIFY?

=> d ibib ab 1-6

L1 ANSWER 1 OF 6 CAPLUS COPYRIGHT 2000 ACS

ACCESSION NUMBER: 1999:495288 CAPLUS

DOCUMENT NUMBER: 131:131512

TITLE: Methods for **separating** a tocotrienol from a

tocol-containing mixture and compositions thereof

INVENTOR(S): Sumner, Charles E., Jr.; Moncier, John D.; Kanel,

Jeffrey S.; Foster, Mary K.

PATENT ASSIGNEE(S): Eastman Chemical Company, USA

SOURCE: PCT Int. Appl., 74 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9938860	A1	19990805	WO 1999-US1571	19990126
W: BR, CN, ID, JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,				
NL,				
PT, SE				
PRIORITY APPLN. INFO.:			US 1998-PV72962	19980129
			US 1998-PV72963	19980129

AB The invention relates to method for **seppg.** tocotrienol from a first tocol admixt. by heating the first tocol admixt. composed of a tocotrienol, at least one **tocopherol**, a fatty acid, and an **esterifying** compd. to **esterify** the fatty acid to produce a second tocol admixt. composed of the tocotrienol, the **tocopherol**, the esterified fatty acid, and the unesterified fatty acid; distg.

the second tocol admixt. with the esterified fatty acid to remove the unesterified fatty acid from the second tocol admixt. to produce a third tocol admixt. composed of the tocotrienol and the **tocopherol**, with substantially removed unesterified fatty acid; distg. the third tocol admixt. for a sufficient time and temp. to substantially remove the tocotrienol and **tocopherol** from the third tocol admixt. to

produce a fourth tocol admixt. composed of the removed tocotrienol, **tocopherol**, and a non-tocol component; and extg. the tocotrienol from the fourth tocol admixt. with an extn. solvent composed of a polar, org. solvent that is miscible with water to produce a two phase system composed of a first phase contg. the majority of the extn. solvent and the second phase, wherein the selectivity of the extn. solvent for tocotrienol with respect to the **tocopherol** is greater than unity, and removing the first phase from the second phase, with the proviso that the extn. solvent is not a neat alc. The invention further relates to compns. produced by the methods of the present invention.

L3 ANSWER 2 OF 6 CAPLUS COPYRIGHT 2000 ACS
 ACCESSION NUMBER: 1999:495287 CAPLUS
 DOCUMENT NUMBER: 131:131511
 TITLE: Methods for **separating** a tocol from a tocol-containing mixture
 INVENTOR(S): Sumner, Charles E., Jr.; Moncier, John D.; Kanel, Jeffrey S.; Foster, Mary K.
 PATENT ASSIGNEE(S): Eastman Chemical Company, USA
 SOURCE: PCT Int. Appl., 73 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9938859	A1	19990805	WO 1999-US1570	19990126
W: BE, CN, ID, JP				
FW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GE, GR, IE, IT, LU, MC,				

PT, SE

PRIORITY APPLN. INFO.:

US 1998-PV72962 19980129
 US 1998-PV72963 19980129

AB The invention relates to a method for **sepg.** tocol from a tocol-contg. admixt. by heating the tocol-contg. admixt. composed of a tocol, a fatty acid, and an **esterifying** compd. to **esterify** the fatty acid to produce a second tocol admixt. composed of the tocol, the esterified fatty acid, and the unesterified fatty acid; distg. the tocol-contg. admixt. with the esterified fatty acid to remove the unesterified fatty acid from the tocol admixt. to produce a tocol admixt. composed of the tocol with substantially removed unesterified fatty acid; distg. the tocol admixt. for a sufficient time and temp. to substantially remove the tocol from the tocol admixt. to produce a tocol admixt. composed of the removed tocol and a non-tocol component; and

extg.

the tocol from the tocol admixt. with an extn. solvent composed of a polar, org. solvent that is miscible with water to produce a two phase system composed of a first phase contg. the majority of the extn. solvent and the second phase, wherein the selectivity of the extn. solvent for tocol with respect to the non-tocol component is greater than unity, and removing the first phase from the second phase, with the proviso that the extn. solvent is not a neat alc.

LE ANSWER 3 OF 6 CAPLUS COPYRIGHT 2000 ACS
ACCESSION NUMBER: 1990:503371 CAPLUS
DOCUMENT NUMBER: 113:103371
TITLE: Recovery of carotenoids, **tocopherols**,
tocotrienols and sterols from esterified palm oil
INVENTOR(S): Goh, Swee Hock; Kam, Toh Seok; Choo, Yen May;
Eng,
Augustine Soon Hock
PATENT ASSIGNEE(S): Institut Penyelidikan Minyak Kelapa Sawit
Malaysia,
Malay.; University of Malaya
SOURCE: Brit. UK Pat. Appl., 21 pp.
CODEN: BAXXDU
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 1
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
GB 2213389	A1	19891129	GB 1988-29427	19881215
GB 2213389	B2	19910904		

PRIORITY APPLN. INFO.: GB 1987-29232 19871215

AB A method for the isolation of the minor nonglyceride components of palm oil or similar vegetable oil contg. free fatty acid and nonglyceride components similar to that of palm oil comprises: (a) **esterifying** the free fatty acid component of the oil with .gtoreq.1 monohydric alcs. to form an esterified oil with a very low free fatty acid content; (b) converting the glycerides into monoesters by transesterification using .gtoreq.1 monohydric alc.; (c) adsorbing the nonglyceride components onto a selective adsorbent to **sep.** them from the esters; and (d) desorbing the nonglyceride components from the adsorbent. The adsorbent is preferably activated alumina, activated C, or silica gel. By the method, carotenes, sterols, **tocopherols** and other nonglyceride components can be isolated. Crude palm oil Me ester was dissolved in MeOH and applied to a C18 reversed-phase column. The esters were eluted

with

MeOH, and then carotenoids were eluted with C₆H₁₄: MeOH (98:2 vol./vol.)
or CHCl₃.

L3 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2000 ACS

ACCESSION NUMBER: 1988:114619 CAPLUS

DOCUMENT NUMBER: 108:114619

TITLE: Concentration of **tocopherols** from soybean sludge by supercritical fluid extraction

AUTHOR(S): Shishikura, Akihiro; Fujimoto, Kenshiro; Kaneda, Takashi; Arai, Kunio; Saito, Shozaburo

CORPORATE SOURCE: Fac. Agric., Tohoku Univ., Sendai, Japan

SOURCE: Yukagaku (1988), 37(1), 8-12

CODEN: YKGKAM; ISSN: 0513-398X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB **Sepr.** of **tocopherols** from soybean sludge (byproduct of soybean oil deodorization) by supercrit. carbon dioxide (SC-CO₂) and nitrous oxide (SC-N₂O) was carried out. This **sepr.** was not possible by supercrit. fluid (SCF) alone and consequently, it was necessary to **esterify** the sludge with EtOH. **Tocopherols** could be concd. by 64%. The **tocopherols** could be recovered by 90% in this fraction by SC-CO₂ extrn. in conjunction with a silica gel column and simple pretreatment. The silica gel column could be regenerated for the most part by extrn. with an entrainer, such as EtOH.

L3 ANSWER 5 OF 6 CEABA COPYRIGHT 2000 DECHEMA

ACCESSION NUMBER: 1988:771818 CEABA

TITLE: Concentration of tocopherols from soybean sludge by

supercritical fluid extraction

Anreicherung von Tocopherolen aus

Sojabohnenrueckstaenden durch superkritische Fluidextraktion

AUTHOR: Shishikura, A.; Fujimoto, K.; Kaneda, T.; Arai, K.;

Saito, S.

SOURCE: Yukagaku (1988) 37(1(392)), p.8-12

CODEN: YKGKAM ISSN: 0513-398X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB **Separation** of **tocopherols** from soybean sludge by supercritical carbon dioxide (SC-CO₂) and nitrous oxide (SC-N₂O) was carried out. This **separation** was not possible by supercritical fluid (SCF) alone and consequently, it was considered necessary to **esterify** the sludge with ethanol. Greater **separation** was possible using sludge esterified free fatty acid with ethanol. **Tocopherols** could be concentrated by more than 64%. The **tocopherols** could be recovered by 90% in this fraction by SC-CO₂ extration in conjunction with a silicic acid column and simple pretreatment. This silicic acid column could be regenerated for the most part by extraction with an entrainer, such as ethanol. (Author)

L3 ANSWER 6 OF 6 JICST-EPlus COPYRIGHT 2000 JST

ACCESSION NUMBER: 880172978 JICST-EPlus
TITLE: Concentration of **tocopherols** from soybean sludge
by supercritical fluid extraction.
AUTHOR: SHISHIKUPA A; FUJIMOTO K; ARAI K; SAITO S
KANEDA T
CORPORATE SOURCE: Tohoku Univ., Sendai-shi, JPN
Koriyama Women's Coll., Koriyama-shi, JPN
SOURCE: Yukagaku (Journal of the Japan Oil Chemists' Society),
(1988) vol. 37, no. 1, pp. 8-12. Journal Code: G0238A
(Fig. 7, Tbl. 2, Ref. 17)
CODEN: YKGKAM; ISSN: 0513-398X
PUB. COUNTRY: Japan
DOCUMENT TYPE: Journal; Article
LANGUAGE: English
STATUS: New

AB **Separation of tocopherols** from soybean sludge by
supercritical carbon dioxide(SC-CO₂) and nitrous oxide(SC-N₂O) was
carried out. This **separation** was not possible by supercritical
fluid(SCF) alone and consequently, it was considered necessary to
esterify the sludge with ethanol. Greater **separation** was
possible using sludge esterified free fatty acid with ethanol.
Tocopherols could be concentrated by more than 64%. The
tocopherols could be recovered by 90% in this fraction by SC-CO₂
extraction in conjunction with a silicic acid column and simple
pretreatment. The silicic acid column could be regenerated for the
most part by extraction with an entrainer, such as ethanol.(author abst.)

=> d his

(FILE 'HOME' ENTERED AT 15:11:03 ON 23 MAR 2000)

FILE 'AGRICOLA, ALUMINIUM, ANABSTR, APILIT, BIOCOMMERCE, BIOTECHNO,
CAEA,
CAOLD, CAPLUS, CBNB, CEABA, CEN, CERAB, CIN, COMPENDEX, CONFSCI,
GENBANK,
INSPEC, INSPHYS, INVESTEXT, IPA, JICST-EPLUS, KKF, KOSMET, METADEX,
NAPRALERT, NIOSHTIC, NTIS, PAPERCHEM2, ...' ENTERED AT 15:11:42 ON 23

MAR

2000

L1 43093 S TOCOPHEROL
L2 2438 S L1 AND SEPARAT?
L3 6 S L2 AND ESTERIFY?

=> s l2 and extract?

18 FILES SEARCHED...
L4 734 L2 AND EXTRACT?

=> s l4 and distil?

27 FILES SEARCHED...
L5 36 L4 AND DISTIL?

=> s l5 and esterif?

L6 7 L5 AND ESTERIF?

=> d ibib ab 1-7

L6 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2000 ACS
ACCESSION NUMBER: 1999:495288 CAPLUS
DOCUMENT NUMBER: 131:131512
TITLE: Methods for **separating** a tocotrienol from a
tocol-containing mixture and compositions thereof
INVENTOR(S): Sumner, Charles E., Jr.; Moncier, John D.; Kanel,
Jeffrey S.; Foster, Mary K.
PATENT ASSIGNEE(S): Eastman Chemical Company, USA
SOURCE: PCT Int. Appl., 74 pp.
CODEN: PIXXD2
DOCUMENT TYPE: Patent
LANGUAGE: English
FAMILY ACC. NUM. COUNT: 2
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9938860	A1	19990805	WO 1999-US1571	19990126

W: BR, CN, ID, JP

RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,

NL,

PT, SE

PRIORITY APPLN. INFO.:

US 1998-PV72962 19980129

US 1998-PV72963 19980129

AB The invention relates to method for **sepg.** tocotrienol from a
first tocol admixt. by heating the first tocol admixt. composed of a
tocotrienol, at least one **tocopherol**, a fatty acid, and an
esterifying compd. to **esterify** the fatty acid to produce
a second tocol admixt. composed of the tocotrienol, the **tocopherol**
, the **esterified** fatty acid, and the unesterified fatty acid;
distg. the second tocol admixt. with the **esterified**
fatty acid to remove the unesterified fatty acid from the second
tocol
admixt. to produce a third tocol admixt. composed of the tocotrienol
and
the **tocopherol**, with substantially removed unesterified fatty
acid; **distg.** the third tocol admixt. for a sufficient time and
temp. to substantially remove the tocotrienol and **tocopherol**
from the third tocol admixt. to produce a fourth tocol admixt.
composed of
the removed tocotrienol, **tocopherol**, and a non-tocol component;
and **extg.** the tocotrienol from the fourth tocol admixt. with an
extn. solvent composed of a polar, org. solvent that is miscible
with water to produce a two phase system composed of a first phase
contg.
the majority of the **extn.** solvent and the second phase, wherein
the selectivity of the **extn.** solvent for tocotrienol with
respect to the **tocopherol** is greater than unity, and removing
the first phase from the second phase, with the proviso that the
extn. solvent is not a neat alc. The invention further relates to
compsn. produced by the methods of the present invention.

L6 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2000 ACS
ACCESSION NUMBER: 1999:495287 CAPLUS

DOCUMENT NUMBER: 131:131511
 TITLE: Methods for **separating** a tocol from a tocol-containing mixture
 INVENTOR(S): Sumner, Charles E., Jr.; Moncier, John D.; Kanel, Jeffrey S.; Foster, Mary K.
 PATENT ASSIGNEE(S): Eastman Chemical Company, USA
 SOURCE: PCT Int. Appl., 73 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9938859	A1	19990805	WO 1999-US1570	19990126
W: BR, CN, ID, JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, PT, SE				

2105
 NL,
 PT, SE
 PRIORITY APPLN. INFO.: US 1998-PV72962, 19980129
 US 1998-PV72963, 19980129

AB The invention relates to a method for **seppg.** tocol from a tocol-contg. admixt. by heating the tocol-contg. admixt. composed of a tocol, a fatty acid, and an **esterifying** compd. to **esterify** the fatty acid to produce a second tocol admixt. composed of the tocol, the **esterified** fatty acid, and the unesterified fatty acid; **distg.** the tocol-contg. admixt. with the **esterified** fatty acid to remove the unesterified fatty acid from the tocol admixt. to produce a tocol admixt. composed of the tocol with substantially removed unesterified fatty acid; **distg.** the tocol admixt. for a sufficient time and temp. to substantially remove the tocol from the tocol admixt. to produce a tocol admixt. composed of the removed tocol and a non-tocol component; and **extg.** the tocol from the tocol admixt. with an **extn.** solvent composed of a polar, org. solvent that is miscible with water to produce a two phase system composed of a first phase contg. the majority of the **extn.** solvent and the second phase, wherein the selectivity of the **extn.** solvent for tocol with respect to the non-tocol component is greater than unity, and removing the first phase from the second phase, with the proviso that the **extn.** solvent is not a neat alc.

L6 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2000 ACS
 ACCESSION NUMBER: 1997:533639 CAPLUS
 DOCUMENT NUMBER: 127:190871
 TITLE: Method of producing (.-.)-tocopherol or (.-.)-tocopheryl acetate
 INVENTOR(S): Jaedicke, Hagen; Grafen, Paul; Laas, Harald
 PATENT ASSIGNEE(S): Basf A.-G., Germany; Jaedicke, Hagen; Grafen, Paul;

SOURCE: Laas, Harald
 PCT Int. Appl., 21 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: German
 FAMILY ACC. NUM. COUNT: 2
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9728151	A1	19970807	WO 1997-EP324	19970124
W: CN, JP, RU, SK, US				
PW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
DE 19603142	A1	19970731	DE 1996-19603142	19960129
DE 19617444	A1	19971106	DE 1996-19617444	19960502
EP 882036	A1	19981209	EP 1997-901601	19970124
F: CH, DE, FR, GB, LI				
PRIORITY APPLN. INFO.:			DE 1996-19603142	19960129
			DE 1996-19617444	19960502
			WO 1997-EP324	19970124

OTHER SOURCE(S): CASREACT 127:190871

AB A method is disclosed of producing (+-)-.alpha.-**tocopherol** or (+-)-.alpha.-tocopheryl acetate by acid catalyzed reaction of 2,3,5-trimethylhydroquinone (TMH) with phytol or isophytol (IP) in a solvent at raised temp. and, where appropriate, subsequent **esterification** of the **tocopherol** thus obtained with acetic anhydride. The method is characterized in that the reaction is carried out in an optionally substituted cyclic five-ring carbonate such as 1,3-propylene carbonate, or in an optionally substituted five-ring lactone such as .gamma.-butyrolactone, as the solvent at temps. of between 50 and 200.degree.. The reaction works particularly well if, following the reaction of TMH and phytol or IP, the **tocopherol** which **separates** out as an upper phase as the reaction mixt. cools and/or the reaction mixt. is **extd.** using a suitable aliph. hydrocarbon, the **tocopherol** is isolated by **distn.** from the **ext.**, and the **sepd.** cyclic carbonate or .gamma.-lactone which can contain excess TMH and acidic catalyst is reused as solvent. In many cases, it has been shown to be advantageous to carry out the reaction of TMH with phytol or IP with removal of the water as azeotrope, using a suitable hydrocarbon and/or in the presence of a mixt. of ortho-boric acid with oxalic acid, tartaric acid or citric acid, or alternatively in the presence of BF₃ etherate as the acidic catalyst. Surprisingly, the five-ring carbonates and five-ring lactones prove to have sufficient stability under the reaction conditions and suitable solvent characteristics to permit the process to be carried out continuously.

ACCESSION NUMBER: 1991:5295 CAPLUS
 DOCUMENT NUMBER: 114:5295
 TITLE: Technology for **separating** vitamin E from the residues of vegetable oil refining
 INVENTOR(S): Qu, Delin; Chen, Peirong; Su, Jianmin
 PATENT ASSIGNEE(S): Tsinghua University, Peop. Rep. China
 SOURCE: Faming Zhuanli Shenqing Gongkai Shuomingshu, 11 PF.
 CODEN: CNXXEV
 DOCUMENT TYPE: Patent
 LANGUAGE: Chinese
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CN 1034367	A	19890802	CN 1987-106187	19870917
CN 1013197	B	19910717		

AB The title named involves: (1) **esterification** of plant (e.g. soybean) residue with EtOH in MeOH at 60-80.degree. for 4 h, (2) washing the reaction product with 60-70.degree. water until the water is neutral, (4) vacuum-**distn.** of the resultant product to give vitamin E, (4) repeating steps of (1) to (3), (5) **extg.** the crude **distillate** with MeOH on MeOH-acetone, (6) allowing to stand to ppt. and removing the ppt., (7) chromatog. of the supernatant on silica gel, and (8) chromatog. on strongly basic cation exchangers (not specified).

L6 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2000 ACS

ACCESSION NUMBER: 1974:567970 CAPLUS
 DOCUMENT NUMBER: 91:167970
 TITLE: Utilization of deodorized condensates from vegetable oils
 AUTHOR(S): Kim, Jum Sik
 CORPORATE SOURCE: Coll. Eng., Hanyang Univ., Seoul, S. Korea
 SOURCE: Taehan Hwahak Hoechi (1974), 18(4), 297-301
 CODEN: DHWHAB
 DOCUMENT TYPE: Journal
 LANGUAGE: Korean

AB The condensates were heated to **sep.** the oil from the water, and the oil was removed. After Me **esterification**, the oil was **distd.** to remove fatty acids. The oil was then saponified and the unsaponifiable matter was **extd.** with hot MeOH and concd. The sterols were crystd. from the concd. soln. The mother liquor was dissolved in MeOH and cooled to -20.degree. to remove residual fatty acids.
Tocopherols were then **sepd.** with an ion exchange resin.

L6 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2000 ACS

ACCESSION NUMBER: 1971:491267 CAPLUS
 DOCUMENT NUMBER: 75:91267
 TITLE: Process report on vitamins. IV
 AUTHOR(S): Ganju, Avtar

CORPORATE SOURCE: India
SOURCE: Indian Chem. J. (1971), 5(10), 40-3
CODEN: ICLJAG
DOCUMENT TYPE: Journal
LANGUAGE: English
AB Five techniques for isolation of **tocopherols** from animal and vegetable fat deodorizer sludge are reviewed: (1) **esterification** of the 20% sterols, 8 **tocopherols**, 20% fatty acid-triglyceride mixt., **sepn.** of the resulting esters by **distn.**, and alc. wash of the residue to remove the sterols; (2) sapon. of the mixt., **sepn.** of unsaponifiable matter (**tocopherols** and sterols), and wash **sepn.**; (3) **sepn.** of **tocopherols** by chloromethylation and redn.; (4) fractional liq.-liq. **extrn.**; (5) ion exchange.

L6 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2000 ACS
ACCESSION NUMBER: 1967:412811 CAPLUS
DOCUMENT NUMBER: 67:12811
TITLE: Utilization of [vegetable oil] concentrate containing

tocopherols. I. Separation of
concentrate components by thin-layer chromatography

and some properties of reducing substances
AUTHOR(S): Takeuchi, Takahiro; Tatsukawa, Toyokazu
CORPORATE SOURCE: Ind. Res. Inst., Kobe, Japan
SOURCE: Yukagaku (1967), 16(4), 185-93
CODEN: YK GKAM

DOCUMENT TYPE: Journal
LANGUAGE: Japanese
AB It is generally accepted that the deodorized condensates obtained in the refining of vegetable oils contain 1-10% **tocopherols**. The condensates are good raw materials for prepn. of .alpha.-**tocopherol** or **tocopherol** concentrates. The concentrate containing 42.6% .alpha.-type **tocopherols** was prepd. by **esterification** and **distn.** of the soybean oil condensate. Thin-layer chromatog. was used to analyze the concentrate components prior to studies on utilization of the concentrate. For analysis of these components, nonpolar and polar developing solvent systems were at least necessary. Hydrocarbons, squalene, and 5 other unknown components were **sepd.** from each other in nonpolar solvent systems, such as naphtha. On the contrary, .alpha., .gamma., and .delta.-**tocopherol**, higher alcs., and sterols were **sepd.**, each giving **sep.** spots in polar solvent systems, such as C6H14-Et2O (7:3 by vol.). These spots could be detected by uv illumination and spraying with solns. such as icidine-C6H6, 50% H2SO4, or Emmerie-Engel reagent. 20 references.

=> log y

COST IN U.S. DOLLARS

SINCE FILE
ENTRY

TOTAL
SESSION

FULL ESTIMATED COST	88.90	89.11
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-6.12	-6.12

STN INTERNATIONAL LOGOFF AT 15:16:59 ON 23 MAR 2000